## LOUISIANA MEDICAID PROGRAM ISSUED: 08/13/24 REPLACED: 09/08/20

#### **CHAPTER 5: PROFESSIONAL SERVICES**

SECTION 5.1: COVERED SERVICES PAGE(S) 2

### **Hyperbaric Oxygen Therapy**

Hyperbaric oxygen therapy is a modality in which the entire body is exposed to oxygen under increased atmospheric pressure.

#### **Covered Conditions**

Reimbursement for hyperbaric oxygen therapy is limited to treatments administered in a hyperbaric oxygen therapy chamber. Hyperbaric oxygen therapy is covered for the following conditions, if deemed medically necessary:

- 1. Acute carbon monoxide intoxication;
- 2. Decompression illness;
- 3. Gas embolism:
- 4. Gas gangrene;
- 5. Acute traumatic peripheral ischemia. Hyperbaric oxygen therapy is a valuable adjunctive treatment to be used in combination with accepted standard therapeutic measures when loss of function, limb, or life is threatened;
- 6. Crush injuries and suturing of severed limbs. Hyperbaric oxygen therapy would be an adjunctive treatment when loss of function, limb, or life is threatened;
- 7. Progressive necrotizing infections (necrotizing fasciitis);
- 8. Acute peripheral arterial insufficiency;
- 9. Preparation and preservation of compromised skin grafts (not for primary management of wounds);
- 10. Chronic refractory osteomyelitis, unresponsive to conventional medical and surgical management;
- 11. Osteoradionecrosis as an adjunct to conventional treatment;
- 12. Soft tissue radionecrosis as an adjunct to conventional treatment;

# LOUISIANA MEDICAID PROGRAM ISSUED: 08/13/24 REPLACED: 09/08/20

CHAPTER 5: PROFESSIONAL SERVICES
SECTION 5.1: COVERED SERVICES

PAGE(S) 2

13. Cyanide poisoning;

- 14. Actinomycosis, only as an adjunct to conventional therapy when the disease process is refractory to antibiotics and surgical treatment; and
- 15. Diabetic wounds of the lower extremities when the wound is classified as Wagner grade 3 or higher and an adequate course of standard wound therapy was not sufficient to lead to healing.